

## Medical History, Timely Physical Exam, & Required Immunizations

Provide signed documentation of a current medical history listing allergies, required medications, and any health conditions or impairments. This should include a report of a physical examination conducted during the 18 months prior to camp as well as a certificate of required immunizations (listed below). This information can normally be obtained with a simple phone call and is commonly referred to by health care providers as a "camp form."

## **Grades Kindergarten – 6**

Kindergarten requirements apply to all students ≥5 years.

DTaP	<b>5 doses;</b> 4 doses are acceptable if the 4 <sup>th</sup> dose is given on or after the 4 <sup>th</sup> birthday. DT is only acceptable with a letter stating a medical contraindication to DTaP.
Polio	<b>4 doses;</b> $4^{th}$ dose must be given on or after the $4^{th}$ birthday and $\geq 6$ months after the previous dose, or a $5^{th}$ dose is required. 3 doses are acceptable if the $3^{rd}$ dose is given on or after the $4^{th}$ birthday and $\geq 6$ months after the previous dose.
Hepatitis B	3 doses; laboratory evidence of immunity acceptable
MMR	<b>2 doses;</b> first dose must be given on or after the $1^{st}$ birthday and the $2^{nd}$ dose must be given $\geq 28$ days after dose 1; laboratory evidence of immunity acceptable
Varicella	<b>2 doses;</b> first dose must be given on or after the 1 <sup>st</sup> birthday and 2 <sup>nd</sup> dose must be given ≥28 days after dose 1; a reliable history of chickenpox* or laboratory evidence of immunity acceptable

## **Grades 7 - 12**

In ungraded classrooms, grade 7 requirements apply to all students ≥12 years.

Tdap	<b>1 dose</b> ; and history of DTaP primary series or age appropriate catch-up vaccination. Tdap given at ≥7 years may be counted, but a dose at age 11-12 is recommended if Tdap was given earlier as part of a catch-up schedule. Td should be given if it has been ≥10 years since Tdap.
Polio	<b>4 doses;</b> $4^{th}$ dose must be given on or after the $4^{th}$ birthday and $\geq 6$ months after the previous dose, or a $5^{th}$ dose is required. 3 doses are acceptable if the $3^{rd}$ dose is given on or after the $4^{th}$ birthday and $\geq 6$ months after the previous dose.
Hepatitis B	3 doses; laboratory evidence of immunity acceptable
MMR	<b>2 doses;</b> first dose must be given on or after the $1^{st}$ birthday and the $2^{nd}$ dose must be given $\geq 28$ days after dose 1; laboratory evidence of immunity acceptable
Varicella	<b>2 doses;</b> first dose must be given on or after the $1^{st}$ birthday and $2^{nd}$ dose must be given $\geq 28$ days after dose 1; a reliable history of chickenpox* or laboratory evidence of immunity acceptable